



Public Works Department,
Park Division
965 Fir Street
(530) 896-7800

Agenda Prepared: 10/9/17
Agenda Posted: 10/9/17
Prior to: 5:00 p.m.

**CITY OF CHICO
BIDWELL PARK AND PLAYGROUND COMMISSION (BPPC)
TREE COMMITTEE**

(Commissioners Hernandez (Chair), Reddemann, Haar)
October 12, 2017, 6:00 p.m.
Municipal Center - 421 Main Street, Conference Room 1

Materials related to an item on this Agenda are available for public inspection in the Park Division Office at 965 Fir Street during normal business hours or online at <http://www.chico.ca.us>.

1. CALL TO ORDER

- 2. REGULAR AGENDA** - All items listed under the Regular Agenda are in the order which is believed are of interest to the public or which require Committee action at this meeting. The items will be considered in the order listed unless the Committee members request a change. Any person may speak on items on the Regular Agenda.

2.1 VOLUNTEER OAK TREE PLANTING AT ONE MILE RECREATIONAL AREA.

Staff will update the Committee on a proposal to plant Oak Trees in Bidwell Park
Recommendation : *This item is for information only*

2.2 CONSIDERATION OF CHICO MUNICIPAL CODE REVISIONS TO CMC 14.40 AND 16.66 AND 16.68

Chico Municipal Code (CMC) Chapter 14.40 entitled "Street Trees" establishes regulations controlling and governing the planting, removal and maintenance of trees and shrubs on city-owned property and right-of-way. CMC Chapter 16.66 entitled "Tree Preservation Regulations" provides for the protection of City and private trees during the development processes. The Committee will continue its review of potential revisions to the CMC Sections.

Recommendation: *Staff recommends that the committee review and provide additional input on staff's comparative study for revision to these Chico Municipal Code (CMC) Sections*

2.3 CONSIDERATION OF TREE EDUCATION AND OUTREACH PROGRAM

At its 8/10/17 meeting, the Committee discussed establishing a public education and outreach program regarding tree care. The Committee will review a draft outline of an outreach program.

Recommendation: *Staff recommends that the committee review and provide input on proposed topics, suggested content and method of delivery of educational outreach.*

3. BUSINESS FROM THE FLOOR

Members of the public may address the Committee at this time on any matter not already listed on the agenda; comments are limited to three minutes. The Committee cannot take any action at this meeting on requests made under this section of the agenda.

4. ADJOURN

Adjourn to the next regular meeting tentatively scheduled for November 9 2017 at 6:00 pm. in Conference Room 1, Chico Municipal Center building (421 Main Street, Chico, California).



Please contact the Park Division Office at (530) 896-7800 if you require an agenda in an alternative format, or if you need to request a disability-related modification or accommodation. This request should be received at least three (3) working days prior to the meeting.



BPPC Tree Committee Report

Meeting Date 10/12/2017

DATE: 10/12/17
TO: Bidwell Park and Playground Commission (BPPC) Tree Committee
FROM: Richie Bamlet, Urban Forest Manager
SUBJECT: Valley oak restoration tree/acorn planting at One Mile Recreation Area, Bidwell Park

RECOMMENDATION:

This report is for information only.

BACKGROUND:

At its 5/16/17 meeting, the City Council authorized Public Works staff to take the lead in pulling a project together for the fall, in coordination with the volunteer groups who are interested in assisting with the planting of acorns at One Mile Recreation Area

Following is an analysis of the Bidwell Park Master Management Plan Vol 1-Draft EIR-April 2007 Appendix C-NATURAL RESOURCES MANAGEMENT PLAN policies regarding Oak tree planting:

3.1.3 OAK MANAGEMENT

3.1.2 OAK WOODLAND MANAGEMENT OBJECTIVES

The oak woodland management program consists of three interrelated objectives:

- ▶ Ensure oak woodland sustainability by increasing recruitment;
- ▶ Protect existing oak woodlands from wildfire; and,
- ▶ Practice responsible oak landscape maintenance.

3.1.3.1 OAK WOODLAND SUSTAINABILITY Many oak stands within Bidwell Park are even-age stands of mature trees with few young trees, saplings, or seedlings. The lack of young oak trees raises questions about the regeneration of these stands. The term “regeneration” refers to the net change in stand structure resulting from the loss of individual trees to mortality and the gain of individual trees through recruitment. Therefore, a lack of recruitment, as is commonly seen in many oak stands within Bidwell Park, is not necessarily indicative of an oak regeneration problem if there is little to no mortality within the stand. Still, the apparent lack of natural oak regeneration at Bidwell Park potentially poses a threat to the long-term viability of its oak woodlands.

The Butte Environmental Council is entering its fourth year of a highly successful oak restoration project focusing on Middle and Upper parks. This effort is funded by the California Wildlife Foundation. It is not anticipated that BEC’s oak restoration program can extend its remit to Lower park. However, Public Works hopes to reach out to BEC and Chico Tree Advocates to benefit from the experience gained in terms of successful project delivery.

Staff met Council member Coolidge 9/25/17 and examined oak restoration areas at Sycamore field, areas devoid of tree cover adjacent and parallel to Woodland avenue and areas around One Mile recreation area.

DISCUSSION.

The prolonged drought has taken its toll on the stands of native oak in the park. From December 2011 to March 2017, the state of California experienced one of the worst droughts on record since records began in 1885. The 2014 drought was considered the worst in 1200 years. As a general rule, it takes a tree a year to recover from a year of drought. By that reckoning, weather conditions would have to create at least average precipitation levels every year through 2023 for trees to fully recover from the recent six-year extreme drought event.

Recent observations in the park have witnessed significant mortality of mature trees. It can be concluded therefore that the combined lack of young seedlings and the accelerated loss of mature tree canopy represents an oak regeneration problem. Trees are not being recruited into the forest canopy faster than they are being lost. Efforts to introduce artificial tree regeneration should be ongoing. With continued efforts, over an extended period an uneven aged canopy will develop. This will produce many benefits such as increased habitat diversity, increased aesthetics and a more resilient forest. There are already successful regeneration attempts near Sycamore field and other places, but more can be done. Efforts to improve the structure and composition of the tree canopy in Bidwell park will also compliment other natural resource practices such as ongoing removal of invasive species in the park.

Researchers Bernhardt and Tedmund (see Attachment A) showed that regeneration is possible from direct seeding of acorns even without irrigation. Browsing from cattle can strongly inhibit natural seeding; but this is not a factor in the Lower park. Vole damage can be a problem, but the addition of mulch and keeping planting spots weed free should reduce losses.

Project implementation:

It is anticipated that Public Works staff will lead the effort to plant new acorns in the Lower park. Staff hope to work in collaboration with volunteers and local non-profits. Public Works will use a modified version of BEC's work plan as a guide (See attachment B). Public Works can provide recycled tree cages for use in areas of the park that have high visitor usage pressure. Where ground is very hard, staff can also assist in planting with a mechanized auger. Any supplemental watering that is deemed necessary can be done by the in-house staff water truck. Incidental costs such as Tubex tree shelters can be obtained at minimal cost using the city in-lieu tree planting fund.

Some of the areas identified, especially the park perimeter along Woodland Avenue may not be conducive to acorn planting. Challenges in successful tree establishment such as ground preparation and ongoing weeding may require that larger saplings are planted. The option of germinating acorns off-site for later transplanting will also be considered. This idea also reopens the idea of re-establishing a city tree nursery.

Public Works staff expect that regeneration efforts in Lower park will be an ongoing phased project spanning many years.

FISCAL IMPACT:

Public Works Director Gustafson indicated that this project could be done by Public Works staff with only a small impact to the Public Works budget.

Reference resource:

http://www.chico.ca.us/document_library/departments/general_services/Parks/Final_Bidwell_Park_Master_Management_Plan_7-08/Final_Bidwell_Park_Master_Management_Plan_6-08/Appendix_C-NRMP.pdf

Attachments:

Attachment A: General Technical Report PSW-GTR 251. A Long-term performance of minimum input oak restoration plantings

Attachment B: BEC Oak restoration project work plan

Long-Term Performance of Minimum-Input Oak Restoration Plantings¹

Elizabeth Bernhardt² and Tedmund J. Swiecki²

Abstract

Starting in 1989, we used minimum-input methods to restore native oaks to parts of their former ranges in Vacaville, California. Each restoration site was analyzed, and only those inputs deemed necessary to overcome expected limiting factors for oak establishment were used. We avoided unnecessary inputs that added to cost and could have unintended negative consequences. All projects were direct-seeded by volunteers using locally collected acorns of valley oak (*Quercus lobata*) and other native oaks. Other inputs included mulch and protection from herbivores (cattle, voles) or mowing crews. Plantings received sporadic maintenance after planting. None of the plantings were irrigated or fertilized. Growth rates and survival show spatial variation at all locations. Multiple project locations now have stands of oaks that have been established at very low cost, validating the minimum input approach. Some very low input plantings had high mortality due to unanticipated impacts from fire and vole outbreaks that greatly exceeded levels previously observed. Lessons learned from the long-term performance of these plantings can be applied in an adaptive management system to accomplish low cost, ecologically sound oak restoration projects in other locations.

Key words: acorns, direct seeding, herbivore protection, interior live oak, *Quercus lobata*, *Quercus wislizeni*, valley oak

Introduction

Valley oak (*Quercus lobata*) was removed from much of its former range to make way for agricultural and urban development, especially in the late 19th and early to mid 20th centuries. Regulatory protections now in place usually require some form of mitigation if valley oaks or other native oaks are removed for development. Restoration plantings in degraded or non-occupied habitat are a common mitigation requirement.

Under optimal soil and water conditions, valley oak is easy to grow, but it is more difficult to establish valley oak in suboptimal sites. In 1989, we began a project to examine low-input methods for restocking valley oaks on formerly forested parcels that were being used for cattle grazing (Bernhardt and Swiecki 1991, 1997). After visiting previous plantings and reviewing the literature, we developed a model to guide the selection of low input cultural methods needed for successful planting at a given site. Demonstration projects established at that time were designed to show whether valley oaks could be established in rangeland settings from acorns without supplemental irrigation. Based on initial success of low input techniques, the model was used to select inputs for additional restoration plantings that occurred from 1993 through 2000 in Vacaville, California. In this paper, we present long-term survival and growth data and discuss the effectiveness of the minimum input techniques employed for Vacaville plantings conducted between 1989 and 2000.

¹ An abbreviated version of this paper was presented at the Seventh California Oak Symposium: Managing Oak Woodlands in a Dynamic World, November 3-6, 2014, Visalia, California.

² Phytosphere Research, 1027 Davis Street, Vacaville CA 95687. (Phytosphere@phytosphere.com).

Methods

All sites were direct seeded using valley oak acorns collected in Vacaville. Acorns were refrigerated in plastic bags between collection and planting. Planting was conducted by volunteers from late October through December after the first soaking rains. Except as noted below, planting sites were prepared by turning over and breaking up the soil with a shovel. Volunteers selected acorns by hand that were free of insect emergence holes, decay, or other obvious defects. At each site, four acorns were planted on their sides at a depth of about 5 cm, spaced 15 cm apart in a square pattern. Planting sites were mulched with wood chips (about 0.8 to 1 m circle) obtained from local arborists, except as noted below. Planting sites were premarked to avoid potentially poor microsites for the 1989, 1993, 1999 and 2000 plantings.

On grazed parcels, planting sites were protected from by cattle browsing with protective Vaca cages (Bernhardt and Swiecki 1991). The cages were 122 cm tall, about 45 cm in diameter, and made of galvanized 12-gauge wire mesh (5 by 10 cm) fencing fabric. Each cage was secured on one side to a T-post and on the opposite side by a 86 cm length of 9.5 to 12.7 mm diameter steel reinforcing bar (rebar) driven into the soil at least 30 cm. As the oaks grew beyond the cages and showed browsing damage, in many instances cages were extended to 180 cm tall by wiring on additional fence fabric. Installation of cage extensions was sometimes delayed, so some trees were suppressed by browsing for several years. Cages were removed after trees grew well above browse height, though cage removal has been delayed beyond this point for many trees.

For areas maintained by mowing, we used 3 m lengths of 16 mm diameter steel reinforcing bar (rebar) bent into U-shaped pins. The straight ends of the rebar were driven into the ground to straddle the planting site, with the top of the pin standing about 1 m above ground level. The pins marked the sites and were intended to deter accidental mowing of the trees and were difficult to vandalize.

Height and survival data were collected in August and September 2014. Height data were collected with the aid of a telescoping measuring pole. Trunk diameter 1.37 m above grade (DBH) was measured with a caliper. Canopy diameter was measured along two perpendicular axes, using either a measuring tape or laser rangefinder. The canopy area was calculated using the formula for an ellipse. Data were analyzed using JMP[®] 9.0.3 statistical software (SAS Inc., Cary NC). Tukey-Kramer HSD was used to separate means following a significant analysis of variance.

1989 planting

The planted area (east and west) was on two generally south facing hillsides divided by a residential street in north Vacaville. The hillsides were commonly grazed for several weeks or longer between December and June. Grazing duration and intensity has varied widely from year to year, and did not occur in some years. At the start of the project, we anticipated browsing by cattle, moisture stress due to weed competition, soil depth and compaction, and vandalism would be the most likely factors to limit restocking success. As part of a study, we tested five treatments, described in detail in Bernhardt and Swiecki 1991. The lowest input treatment had no protection from cattle grazing. The 2014 assessments did not include relocating the planting sites for this treatment; almost all of these were vacant after the first few years. The other four treatments were protected by Vaca cages. These treatments included augering with a 10-cm diameter bit to a depth of 45 to 60 cm and mulching with landscape fabric covered with woodchips; landscape fabric and wood chip mulch without augering; and preselection of planting sites based on penetrability by a

ATTACHMENT A

Proceedings of the 7th California Oak Symposium: Managing Oak Woodlands in a Dynamic World

steel probe (30 cm vs. 45 to 60 cm depth) with only dry grass mulch. We planted 30 sites per treatment per hillside. The planted area was about 1.4 ha on the east side and 1.2 ha on the west side.

1993 planting

The planted area was 2.1 ha on hill slopes and wide drainages along a watershed in a cattle-grazed area in Lagoon Valley. Soils were relatively deep and loamy, with some areas of slippage near the main drainage. Grazing typically began in late December and continued into late April or May. We anticipated that browsing by cattle and rodent chewing would be limiting factors at this location. In addition to valley oak, interior live oak (*Quercus wislizeni*) and California buckeye (*Aesculus californica*) were direct-seeded at this location. We planted 113 sites, all of which were protected with Vaca cages and mulched with wood chips. We used 61 cm tall, 15 cm diameter aluminum window screen cylinders buried to a depth of 15 cm for 82 of the planting sites to protect against rodent herbivory. Screens were folded closed at the top and open below ground. Screens were opened as soon as the plant inside reached the top of the cylinder, and were eventually removed. Empty planting sites were replanted through 1996.

1999 planting

This planting consisted of 94 planting sites over 1.2 ha in a nearly level valley floor area. Sites were along the west side of a 0.9 km long section of abandoned railroad right of way, bordered by residential and commercial development. We anticipated vandalism and accidental mowing of seedlings would be limiting factors. Sites were mulched with wood chips and rebar U-pins were installed after planting, but before seedling emergence.

2000 planting

This planting consisted of 98 planting sites over 1.2 ha along the east side of the same right of way described above for the 1999 planting. We anticipated vandalism, accidental mowing, and soil compaction would be limiting factors. A paved biking trail was constructed on the railroad bed after the 1999 planting but before the 2000 planting. Soil on the east side appeared to be more compacted than on the west side. To mitigate surface compaction, prior to planting, each premarked planting site was excavated to a depth of about 60 cm and refilled using a backhoe. Rebar U-pins were installed prior to planting. Otherwise, sites were planted and maintained as described for the 1999 planting.

Other plantings

Other plantings took place in 1994, 1996, 1997, and 1998 on hillside open-space in Lagoon Valley that was formerly grazed. Planting sites were not premarked or protected in any way and no followup maintenance was conducted. We anticipated that rodents and weed competition for soil moisture would be limiting factors. Sites were mulched with wood chips.

Calculation of canopy cover

For each location, the canopy area of trees with DBH >0 was summed and divided by the total area of the planting. Canopy spread was not recorded for some trees; for those, canopy area was calculated from DBH using the formula based on a regression (fig. 2). Canopy spread of existing mature trees was measured from digital images using Google Earth[®].

Results

Survival

Emergence and survival for valley oaks was initially high (> 95 percent of planting sites) for all planting years. Surviving tree densities for locations shown in table 1 vary from 37 to 77 trees per ha. In the oldest (1989) plantings, survival of planting sites protected by Vaca cages (70 percent west, 74 percent east) did not change between the 2014 and 1995 censuses (Bernhardt and Swiecki 1997). No significant differences in survival were seen between the four planting treatments in Vaca cages. We did not attempt to relocate the unprotected planting treatment, but 13 unprotected oaks, all less than 1 m tall, were observed. Most of these were originally planted in Vaca cages that were subsequently lost due to vandalism; no more than two or three may be survivors from the unprotected treatment.

Table 1—Survival of valley oaks, and percent single, double, or multi-trunked at five planting locations

| Planting | Age in 2014, years | Percent | | | | |
|------------------------|-----------------------|----------|-------|-----------|------------|-----------------|
| | | survival | DBH>0 | one trunk | two trunks | multiple trunks |
| 1989-east ^a | 25 | 74 | 72 | 60 | 31 | 9 |
| 1989-west ^a | 25 | 70 | 58 | 20 | 10 | 70 |
| 1993 | 21 | 84 | 69 | 64 | 28 | 8 |
| 1999 | 15 | 91 | 83 | 40 | 44 | 16 |
| 2000 | 14 | 94 | 84 | 30 | 43 | 27 |

^aSites in Vaca cages only.

In May 2007, a grass fire burned most of the 1989 west planting, affecting 77 percent of the planting sites. Trees in the burned area were either topkilled and have since resprouted from ground level (65 percent of all surviving trees) or were at the edge of the burned area and scorched but not topkilled (7 percent of all surviving trees). The remainder appear unaffected by the fire. Due to prolific sprouting of topkilled trees, fire did not decrease survival compared with 1995 census data.

Survival of the 1993 planting (table 1) was affected by both cattle and voles (presumably *Microtus californicus*). Within the first year of planting, four sites had to be abandoned because cattle had pushed over and mangled the Vaca cages. Because of the heavier cattle use at this location, we reinforced cages by wiring them to the T-posts and adding additional T-posts and rebar to the most impacted sites. Wire screens afforded very short-lived protection; within a year or two, we had to open the screen cylinders because oaks and buckeyes had reached the closed tops. This planting was affected by a surge in the local vole population in 1999. Vole populations and associated damage attained levels we had not previously seen locally. Scattered vole damage have been observed sporadically in the planting after

ATTACHMENT A

Proceedings of the 7th California Oak Symposium: Managing Oak Woodlands in a Dynamic World

1999. To date, 16 valley oaks with DBH values ranging up to 12 cm have been girdled and topkilled by voles, and other trees have been partially girdled. Survival of the 15 sites planted with California buckeyes is 100 percent. Eight of 18 sites planted with interior live oak acorns contain surviving trees.

Survival was highest overall at the two youngest plantings (table 1). The 1999 and 2000 plantings are mowed once or twice annually by city crews in May and late summer. In early May 2000, 46 sites in the 1999 planting were mowed by inmate crews using string trimmers while the city coordinator for the site was on vacation. Plants were mowed to heights as low as 5 cm. The mowed seedlings resprouted, and although they were significantly shorter than the unmowed seedlings by August 1999 (8.5 cm vs 15.7 cm), survival on a planting site basis (at least one seedling per site) did not differ. Accidental mowing of marked sites with rebar pins has been an ongoing problem for trees that are shorter than 1 m. Some rebar pins have been rolled over and destroyed by mowing equipment and some sites that have not had the pins replaced (six in 1999 planting, seven in 2000 planting) have been mowed down annually.

Unprotected and unmaintained planting sites on nongrazed grassy upland sites showed good initial emergence but had low long-term survival. For example, the 1994 hillside planting had 95 percent initial survival of valley oak sites the August after planting. By 3 years after planting, survival of valley oak planting sites was 57 percent. In 1999, 5 years after planting, high vole populations in the Lagoon Valley area caused severe damage to this and other unprotected plantings (1996, 1997, 1998), as well as established natural regeneration in nongrazed areas. Some established natural oak saplings with basal diameters greater than 7 cm were killed by girdling of the stem above and below the soil line. Less intense vole population outbreaks have occurred at several times since 1999, mainly in grassy, nongrazed sites. In addition to vole herbivory, large portions of the 1996, 1997, and 1998 plantings were burned in several successive years. Consequently, only a few surviving valley oaks remain from these plantings. Most of the survivors are in mid- to lower-slope positions where oaks unaffected by voles have grown well. The maximum DBH among these survivors is 35 cm. Although deer frequent the area, we have not observed height growth being suppressed by deer browsing at these sites.

Growth

Tree size parameters showed relatively wide distributions for all plantings (table 2, fig. 1). The largest trees were in the oldest plantings (fig. 1) but all sites contained small trees that were less than 1.4 m tall and had no DBH. At each location, trees of similar sizes tended to be spatially clustered, indicating the strong role of underlying soil conditions, such as depth and available moisture, as primary determinants of tree size. Tree DBH, height, and canopy spread in the different plantings overlapped substantially (figs. 1, 2) despite the difference in ages for the plantings. Across all plantings, DBH was highly correlated with height and canopy spread (fig. 2).

Table 2—Comparison of size attributes (mean \pm standard deviation), and calculated canopy cover of valley oaks at 5 planting locations. Means followed by the same letters are not statistically different from one another according to Tukey-Kramer HSD

| Planting | DBH, cm | Height, m | Canopy spread, m ² | Canopy cover, pct |
|------------------------|--------------------------------|-----------------------------|-------------------------------|-------------------|
| 1989-east ^a | 13.2a \pm 8.6 | 5.0a \pm 3.2 | 12.1b \pm 13.9 | 10.0 |
| 1989-west ^a | 11.9ab \pm 6.6 ^{b2} | 4.5a \pm 2.1 ^b | 13.5b \pm 9.0 ^b | 3.6 |
| 1993 | 14.3a \pm 10.8 | 5.3a \pm 3.1 | 31.2a \pm 25 | 10.5 ^c |
| 1999 | 11.0ab \pm 6.7 | 5.4a \pm 2.6 | 15.6b \pm 12.6 | 9.6 |
| 2000 | 8.7b \pm 6.0 | 4.8a \pm 2.6 | 11.4b \pm 9.9 | 7.0 |

^aSites in Vaca cages only.

^bUnburned trees only.

^cIncludes canopy contributions of interior live oaks and protected natural trees.




Figure 1—Distribution of DBH values for plantings shown in tables 1 and 2. Dark shading for second chart from left represents topkilled trees in the burned section of the planting.

In the 1989-east planting, the largest trees were located at the bottom of the slope along an alluvial fan. Vaca cages have been removed from these large trees, so initial treatment assignments, which were marked on the T-posts, were not available. However, all initial treatments were represented among these uniformly large trees. About one third of the remaining trees in the 1989-east had treatment codes still visible. For these trees, the original treatment did not significantly affect DBH, height, or canopy spread.

ATTACHMENT A




Figure 2—Relationships between DBH and tree height (left) and DBH and canopy area (right) across all plantings. Planting year symbols: 1989-east = o, 1989-west = +, 1993 = \diamond , 1999 = X, 2000 = Δ . Regression equations: Left, Height, $m = 1.96 + 0.287 \times \text{DBH cm}$, $R^2 = 0.795$, F ratio $P < 0.0001$. Right: Canopy area, $m^2 = -6.174 + 1.502 \times \text{DBH cm} + 0.0529 \times (\text{DBH cm} - 12.32)^2$, $R^2 = 0.772$, F ratio $P < 0.0001$.

Unlike the 1989-east planting, the largest trees in the 1989-west planting were along the hill crest and on a southeast-facing slope. In the 1993 planting, the largest trees were mostly on a north-facing slope and in broad drainages, and were more common in the lower elevations of the planting, which spanned about 85 m of elevation change. Spatial clustering of tree sizes was also evident on the level planting sites used for the 1999 and 2000 plantings, with poorer growth occurring in the more compacted and poorly-drained soils near the center portion of the linear area.

All 55 trees topkilled by the 2007 fire in the 1989-west planting resprouted prolifically. Nine of these had not yet attained a DBH. For the remaining resprouts, the mean DBH of the largest live stem was 3.9 cm (range 0.6 to 10.1 cm, $n = 46$), significantly smaller (t test $P < 0.0001$) than the DBH of trees outside the burned area shown in Table 2. We were able to measure DBH of the fire-killed trunk at 42 of the burned sites, these averaged 4.2 cm (range 1 to 10.8 cm). The original planting treatments did not significantly affect DBH, canopy area, or height of either nonburned or resprouted topkilled trees in the 1989-west planting. The non-burned trees in the 1989-west planting did not differ significantly from trees in the 1989-east planting with respect to DBH, height, or canopy area (table 2).

As part of the 1993 planting, we installed Vaca cages around browse-suppressed (less than 1 m tall) natural regeneration that was near a mature valley oak in the lower portion of the planting area. We protected six valley oaks that were well beyond the canopy of the existing oak, leaving some adjacent and nearby oaks unprotected. These oaks grew rapidly when protected from winter-spring cattle browsing and most required extended cages within a few years. In 2014, DBH of these trees ranged from 17 to 49 cm. Unprotected regeneration in the same area has remained less than 1 m tall.

Among the interior live oaks in the 1993 planting, seven trees had DBH values ranging from 4.2 to 47 cm, heights from 2.6 to 9 m, and canopy spreads from 3 to 98 m^2 . One interior live oak tree had been topkilled by voles and had resprouted prolifically. Its largest sprout was 1.5 m tall with a DBH of 3 mm.

At most planting spots more than one of the four planted acorns emerged, but we did not thin multiple seedlings. By 2014, the number of planting spots with multiple

oaks varied from location to location (table 1). We excluded the burned, topkilled plants from further analysis. There was no correlation between the DBH of the largest stem at each planting spot and the number of stems at each spot. One seedling usually became dominant and the additional seedlings were suppressed. Among multiple-tree sites where at least one tree had attained a DBH, 54 percent had a second tree that was at least 1.37 m tall. The average DBH of the largest stem was significantly greater than the average of the second stem that was at least 1.37 m tall. Codominant stems were most common in the 1999 and 2000 plantings. The trunks of these trees appear to fuse at the base where they come in contact, though it is not clear whether the appressed trunks actually become grafted together or remain separated by included bark.

The larger oaks at all planting sites have been producing acorns for a number of years. Some small seedlings arising from these were observed in the 1989 and 1993 plantings, although they have been suppressed by grazing.

Removing protective hardware

Removal of protective Vaca cages from large trees was generally delayed beyond the optimum timing and has only been completed to the degree needed for the 1989-east and 1993 plantings in the past 4 years. In a number of instances, especially where the height of the Vaca cage had been extended, branches had grown through the wire mesh. Branches and trunks generally grow around the wire, which can become completely embedded in the tree and can only be cut away at the bark surface. In some cases, straight sections of wire could be cut off and pulled out of the stems, but removal of the Vaca cage at this late stage was time consuming and sometimes difficult. Rebar used to anchor the cages also became embedded at the base of the largest tree trunks and had to be cut off in some instances. Similarly, tops of T-posts were cut off at or below grade with a portable reciprocating saw in instances where roots had grown over the anchor plates and the posts could not be pulled. Rebar U-pins have not yet been removed from most of the 1999 and 2000 plantings. Removal of the pins is overdue for many trees that have begun to grow around rebar that is against the trunk.

Discussion

Results from these plantings show that restoration of valley oaks from direct seeded acorns without irrigation can be successful, even in rangeland settings. These observations confirm earlier results showing that cattle strongly inhibit natural or artificial valley oak regeneration by browsing seedlings and saplings (Bernhardt and Swiecki 1997). No more than a few of the 60 original unprotected sites from the 1989 plantings have survived, and these and other nonprotected seedlings at this lightly grazed site are less than 1 m tall. Natural browse-suppressed oak regeneration that was protected from grazing in the 1993 planting grew as well as the planted acorns, whereas short unprotected oaks also failed to grow above 1 m. In contrast, the few surviving oaks in the unprotected plantings in nongrazed areas nearby have grown into small trees, even though deer are common in this area. Protecting browse-suppressed natural regeneration in cattle-grazed areas can be used to recruit valley oak (Bernhardt and Swiecki 1997) and blue oak (McCreary and others 2011), but opportunities are limited where mature trees are very sparse or absent.

By controlling the growth of herbaceous vegetation, grazing can indirectly promote growth and survival of protected oak seedlings by making the habitat less

ATTACHMENT A

favorable for voles (Bernhardt and Swiecki 1997, McCreary 2001, McCreary and others 2011). Valley oak seedlings planted in nongrazed grassy areas suffered much more attrition from vole herbivory than seedlings in the nearby grazed parcel, although relatively large saplings were damaged even in the grazed parcel. Although voles have been reported to seriously limit oak survival and growth at other sites (Tecklin and McCreary 1993), vole herbivory was not a problem at all Vacaville locations. Significant damage from voles or other rodents has not been observed in the 1989 plantings, which appear to have habitat suitable for voles, or in the 1999 and 2000 urban plantings where habitat is not suitable for voles. Because vole damage can occur when trees are well beyond the size that shelters or screens are useful, habitat modification and other population control measures may be worthwhile inputs.

Although Vaca cages, U-pins, and other protective devices may be necessary to recruit trees, additional labor and cost is required to remove these devices. Timely removal can minimize the work required and allow for easier reuse of materials while avoiding potential damage to trees. Monitoring and an available source of labor are needed to optimize the removal of cages and other devices. Because oaks can grow at widely different rates, hardware removal may need to extend over many years.

The spring grass fire that burned the 1989-west 18 years after the planting did not affect survival of this planting, but the topkill of trees by fire set the growth of these trees back many years. In contrast, repeated grass fires in much younger plantings (1 to several years old) along with vole activity resulted in high attrition rates in several unmaintained plantings in nongrazed areas. We previously showed strong negative effects of fire on growth and survival of natural blue oak seedlings in this same general area (Swiecki and Bernhardt 2002).

Because continued tree attrition due to fire, vole damage, or other factors can continue for many years after initial establishment, we have avoided thinning planting sites to a single tree per site. The presence of multiple seedlings per site has not shown a negative effect on tree growth up to 25 years after planting. Similar results have been seen by Tyler and Moritz (*Quercus lobata* seedlings and conspecific neighbors: Competitors or allies? these proceedings). In most sites, a single tree becomes dominant even though suppressed additional trees persist for many years. However, these suppressed trees may still be capable of being released if the dominant tree is killed or severely damaged.

Canopy cover after 14 to 25 years was still relatively low at all locations (table 2). Nevertheless, projections based on likely mature canopy spreads show current tree densities are adequate to result in moderate to high canopy cover when trees reach mature sizes. For example, the average canopy area of mature oaks near the 1993 planting site is about 250 m². If all of the surviving valley and interior live oaks in this planting (37 oaks/ha, lowest density of the sites) attained typical mature canopy spread, canopy cover would be about 93 percent. This argues against planting at initially dense rates to increase canopy cover in the short term and instead suggests a strategy of phased planting over time if needed to increase cover. Furthermore, because soil variations strongly influence tree growth over the long term, increasing density in areas with inherently slow growth is a poor strategy for increasing canopy cover.


Acknowledgments

We thank the many volunteers who have assisted with data collection, tree planting, or cage removal over the years, most recently Steve Edmundson, Daniel Garcia, Pam Muick, and

Rhett Richardson. We also thank the City of Vacaville for support and materials and the California Department of Forestry and Fire Protection for initial funding.

References

- Bernhardt, Elizabeth A.; Swiecki, Tedmund J. 1991. **Minimum input techniques for valley oak restocking.** In: Standiford, Richard B., tech. coord. Proceedings of the symposium on oak woodlands and hardwood rangeland management. Gen. Tech. Rep. PSW-126. Berkeley, CA: 2–89.
- Bernhardt, E.A.; Swiecki, T.J. 1997. **Effects of cultural inputs on survival and growth of direct seeded and naturally occurring valley oak seedlings on hardwood rangeland.** In: Pillsbury, Norman H.; Verner, Jared; Tietje, William D., tech. coords. Proceedings of the symposium on oak woodlands: ecology, management, and urban interface issues. Gen. Tech. Rep. PSW-160. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 301–311.
- McCreary, D.D., 2001. **Regenerating rangeland oaks in California.** Oakland, CA: University of California Agriculture and Natural Resources Communication Services Publication 21601. 71 p.
- McCreary, D.; Tietje, W.; Davy, J.; Larsen, R.; Doran, M.; Flavell, D.; Garcia, E. 2011. **Tree shelters and weed control enhance growth and survival of natural blue oak seedlings.** California Agriculture 65: 192–196.
- Swiecki, T.J.; Bernhardt, E.A. 2002. **Effects of fire on naturally occurring blue oak (*Quercus douglasii*) saplings.** In: Standiford, Richard B.; McCreary, Douglas; Purcell, Kathryn L.; tech. coords. Proceedings of the fifth symposium on oak woodlands: oaks in California's changing landscape. Gen. Tech. Rep. PSW-GTR-184. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 251–259.
- Tecklin, J.; McCreary, D.D. 1993. **Dense vegetation may encourage vole damage in young oak plantings.** Restoration and Management Notes 11: 153.



Oak Woodland Restoration in Middle and Upper Bidwell Park
 Butte Environmental Council
www.becnet.org
 Length of project: 2014-2018

Problem Statement

Only a third of California’s oak woodlands remain, and in Butte County both Valley and Blue Oaks are not regenerating adequately to ensure ecological survival. There is also a great need for those who use Bidwell Park to better understand what healthy ecological function looks like, as well as become invested in the park’s restoration.

Background

In the spring of 2014 BEC was awarded \$35,000 from the California Wildlife Foundation to implement an oak planting project on publically protected wildlands. BEC staff met with Dan Efseaff and Ruben Martinez in early March to determine the potential for this project to be implemented in upper Bidwell Park. With important feedback and the go-ahead to develop a proposal for the Bidwell Park and Playground Commission, BEC developed the following work plan.

The goals of this effort are to restore oak woodlands in Middle and Upper Bidwell Park, to protect existing oak seedlings, and to engage park users, community members, and students in the tree planting effort. The work plan was developed based on the feedback from numerous local stakeholders and experts who have experience with oak planting efforts, from publications by the California Oaks Foundation, by reviewing documents including the Butte County Oak Woodlands Management Plan and the Bidwell Park Master Management Plan, and through interviews with Magic Inc. Releaf in Palo Alto.

This project supports the goals of the Bidwell Park Master Management Plan including; involvement of the public in oak woodlands restoration; increasing oak recruitment; and supplemental oak planting.

Timeline

| | | |
|------|--------------------------------------|---|
| 2014 | Fall (September/October) | Collect and store acorns |
| | After first rains (October/November) | Four community planting days Map planting locations |
| 2015 | April | Two community maintenance days Protect existing seedlings Install enclosure caging around two small areas of drip line seedlings. |
| | May through September | Weekly watering |
| | Fall | Collect and store acorns |
| | After first rains (October November) | Four community planting days Map planting locations |
| 2016 | April | Two community maintenance days Protect existing seedlings |
| | May through September | Weekly watering |
| 2017 | April | Oak Assessment Day |
| | May through September | Weekly watering |

Goals and Strategies

Goal 1: To restore oak woodlands in Upper Bidwell Park.

Strategy:

1. Apply for a permit to collect Valley and Blue oak acorns in Bidwell Park.
2. Collect, store, and clean acorns for planting.
3. Purchase and collect supplies and materials.
4. Plant 100 sites the first year and 100 sites the second year for a total of 200 trees.
Depending upon the Valley Oak (*Quercus lobata*) and Blue Oak (*Quercus douglasii*) acorn crop, plantings will be from acorns. The backup plan will be to use seedlings from Floral Native Nursery. Second year locations to be determined upon further discussion with park staff.

Goal 2: To protect existing oak seedlings.

Strategy:

1. Identify and map 25 seedlings to protect from browsing, near plantings. Those protected will be a mix of city planted trees that have outgrown their cages as well as wild trees.
2. Install/replace caging, clear weeds, and apply mulch in the spring of 2015.
3. Install enclosure caging around two small areas of drip line seedlings located across the road from parking lot G (Alligator Hole) to encourage vertical growth, and to research the effects of browsing on regeneration.

Goal 3: Engage park users, community members, and students in the tree planting effort.

Strategy:

1. Develop educational and involvement outreach for the park kiosks; including on-site tabling, jug distribution, and contacting groups.

2. Organize four planting and cage installation days in the 2014 fall and winter.
3. Form core team of tree people, and calendar for group participation.
4. Develop a way for trail users to participate individually by bringing an extra gallon for the trees, and create a sense of 'adoption' within various park user groups.
5. Collaborate with interested 6th through 12th grade teachers to develop research projects and guidelines for students.

Locations – Map and photos below

1. North Rim Trail parking lot B and near cross (total plantings – 25)
2. Chico Rod and Gun Club parking lot / south side of Horseshoe Lake (total plantings – 25)
3. Main parking lot E (total plantings – 30)
4. Bidwell Golf Course (total plantings – 20)

Method for acorn and seedling planting

1. Clear a section 2 ft. in diameter around planting location of grass and weeds.
2. Dig a ~ 8 inch hole.
3. Bury a screen cylinder ~6 inches deep with a few inches or so above ground.
4. Plant the oak seedling or acorn, making sure to compact the soil around the roots.
5. Place Tubex over the seedling or acorns and secure with a wooden stake.
6. Place newspaper, rocks, and wood chip mulch around the seedling.
7. Water.
8. Install bird protection at the top of the Tubex.
9. Explore options for labeling and mapping each tree location.

Method for spring maintenance and summer watering

1. From 8:30am to 11:00 am each Saturday or Sunday May through September (~20 weeks) water seedlings.
2. Fill two 55 gallon barrels loaded onto a truck or trailer at the Rod and Gun Club, use gallon containers and buckets to water.
3. Investigate storage options for tools and supplies at the Rod and Gun Club or Golf Course.
4. Weed as needed and depending upon group size.

Method for protecting existing seedlings

1. Clear a three foot diameter bare area around the seedling.
2. Install newspaper and apply wood chips.
3. Using wooden posts and utility fencing form an enclosure around the seedling to protect against deer browsing.
4. Revisit the site to maintain enclosure and weed.

Maintenance

1. Plant for the first two years, maintaining and water all years.
2. At the projects end, BEC will produce a final report.
3. Research funding opportunities to continue the effort.
4. There are no expectations for city maintenance.

Planting Locations

#1 North Rim trail parking lot B and near the cross.



#2 Rod and Gun Club parking lot and south side of Horseshoe Lake.




#3 Main parking lot E.



#4 Bidwell Golf Course.







DATE: 10/12/17
TO: Bidwell Park and Playground Commission (BPPC) Tree Committee
FROM: Richie Bamlet, Urban Forest Manager
SUBJECT: Consideration of Revisions to Chico Municipal Codes Chapters 14.40 and 16.66 AND 16.68

RECOMMENDATION

Staff recommends that the Committee continues its review and provide input on Staff's proposed revisions and/or provide other revisions to Chico Municipal Code (CMC) Chapters 14.40 and 16.66 and 16.68.

BACKGROUND

At its last meeting in August, the Committee began its review of the following Chico Municipal Code Sections related to Trees:

- CMC Chapter 14.40 entitled "Street Trees" establishes regulations controlling and governing the planting, removal and maintenance of trees and shrubs on city-owned property and right-of-way.
- CMC Chapter 16.66 entitled "Tree Preservation Regulations" provides for the protection of City and private trees during the development processes.
- CMC Chapter 16.68 is entitled "Voluntary Tree Heritage Program" The purpose and intent is to identify and promote public awareness of, maintain, and protect designated trees within the city. A copy of this sections is Attached as Attachment A.

DISCUSSION:

A. HERITAGE TREE APPLICATION FEE.

In February 2010, the Chico Municipal Code was amended to establish a Voluntary Heritage Tree Program (Section 16.68.010).

Examination of the history of development of the Heritage tree ordinance shows that the application fee was once reduced. City Council Agenda Report 11/2/10 outlined a case to reduce the fee from \$307 to \$150. The reasons cited for the reduced fee where a) staff processing time was only approximately two to three hours. The \$307 fee was deemed to be needlessly high and was deterring potential applicants from submitting Heritage tree nominations. Since the time of the price reduction, there have been few applications. Many citizens have cited the cost to be prohibitive. The fee is currently \$168.

1. There is a need to reinvigorate the voluntary Heritage tree program. Since the time of inception, some of the nominated trees have blown down and no new trees have been nominated. Therefore, it is proposed to reduce the Heritage Tree application fee to zero. A survey of cities in California revealed that no other cities levy a fee for voluntary heritage tree nominations.
2. The UFM presents a plan that could result in a cost-neutral Heritage tree program that is free to citizens. It is proposed to develop protocols that will allow Heritage tree applications to be processed in an efficient and effective manner. It is proposed that applications can be accepted year-round but will only be processed and submitted to BPPC committee and City Council for approval on an annual basis at a time to be determined.
3. Depending on the volume of applications an upper limit can be placed on the number of nominations that will be considered each year. It is anticipated that the current practice of installing a plaque could become a volunteer project. Non-profits, donations or an Eagle scout project could manage that part of the Heritage tree nomination process. Utilizing a cost-effective sign to be attached to the tree instead

or mounting on a plinth could be investigated also. Simple tree botanical plaques are currently \$18 and can be ordered online.

If a change to the permit fee for the voluntary Heritage tree program is approved, a program of public education, outreach and promotion of the program would be instigated before the CMC changes were enacted.

B. STREET TREE PERMIT FEE AND SPHERE OF REGULATION.

1. Permit Fee

In tandem with the proposal to make Heritage tree applications free, it is proposed to explore introducing a fee for applications to prune or remove a City street tree. The City already levies a fee on tree permits tied to development. A survey of California cities revealed that many cities charge a fee to cover processing costs for such street tree permits. Planting permits should remain free.

Table 2 summarizes fees charged for work to city trees by some neighboring cities. Examination of the table shows that there is a split between free permits and permits charging a small fee.

2. Homeowner Pruning

It is also proposed to introduce a program whereby *de minimis* incidental pruning on city trees is permissible without the need for a formal permit at all. This would have the benefit of releasing staff from the burden of a large backlog of service request for minor work. This would allow residents to remedy minor tree issues adjacent to their properties without cost or administrative burden or fear of civil or administrative action against them. Residents would have the option to employ a paid gardener or landscaper to do the same minor tree work if they chose to do so.

It is noted that some cities operate a similar scheme without any major issues as follows:

City of Sacramento

The City of Sacramento in its recently overhauled tree ordinance categorizes tree work into two categories, "Regulated work" and "Routine work"

Exert from Sacramento City Ordinance 2016-0026:

"Routine maintenance" means minor pruning*; irrigation; mulch application, mowing or trimming grass or other ground cover close to a tree; application of fertilizer, insecticides, or herbicides in accordance with their label; or any other similar acts that promote the life, growth, or health of trees. Any procedure, technique, or practice that is expressly prohibited under the current ANSI A300 standards, including topping, is not routine maintenance.

*"Minor pruning" means the removal of dead branches; or cutting of roots or branches less than two inches in diameter, measured at the location of the cut, from a private protected tree in a cumulative amount of no more than 10% of a combination of the root system and tree crown within a twelve-month period.

The essence of the code language is that residents are permitted to do minor work on private trees that otherwise meet the criteria to be regulated under the city code.

The City of Fresno

The City of Fresno takes deregulation of minor work a step further and operates a program whereby residents can do minor tree work on City trees without a permit. Notes from discussion with Dan Turner, Forestry Supervisor, City of Fresno 10/6/17:

The City of Fresno allows minor pruning on city owned trees by residents without permit. Work is allowed with the following limitations:

- Residents feet must be kept on the ground; no ladders or ropes allowed. Work can be done with hand tools only. No chainsaws. Maximum 1-inch diameter branches only.
- Typical scenarios where residents utilize this option; pruning around mailboxes, clearing sidewalks and driveways and roof lines of minor encroachment.
- Mr. Turner indicated that there have been no liability issues with this program in the nine years he has worked for the City of Fresno.
- The proposed City of Chico permit structure could be broken down into two broad categories. Minor tree work incurs no paperwork or cost. Major work incurs paperwork for a fee.

Table 1: Summary of proposed regulated and unregulated tree work on city trees located in the public right of way.

| Work type | Permit structure | Tools required | Who does the work? |
|--------------------------------------|--------------------|--|-------------------------------------|
| Minor trimming <2" branches or roots | No permit required | Simple hand tools | Homeowner or gardener/landscaper |
| Major pruning>2" Tree Removals | Permit required | Professional equipment and power tools | Approved professional arborist only |

Table 2: Fee comparison for Heritage tree applications and permit requests for work on city trees.

| | CHICO | SACRAMENTO | DAVIS | SANTA ROSA | REDDING | SAN JOSE | SAN LUIS OBISPO | WEST SAC | SAC COUNTY |
|---------------------------------|----------|---|---------|------------|---------|----------|-------------------|----------|------------|
| HERITAGE TREE APPLICATION | \$150.00 | FREE | FREE | FREE | Free | FREE | FREE | FREE | NA |
| FEE FOR TREE PERMIT (CITY TREE) | FREE | 50/100/110/200 REMOVAL/ROOT/PLANTING/PLANTING | \$54.00 | FREE | Free | FREE | \$81 IF PROCESSED | \$25.00 | \$31.50 |

Below are links to references pertinent to the topics discussed above. If a change to the the permit fee and regulation structure are approved, a program of public education and outreach would be instigated before the CMC changes were enacted.

Attachments:

Attachment A: CMC 16.66 – Heritage Trees

Attachment B: (See reference links below)

Reference resource links:

http://www.isa-arbor.com/education/resources/educ_treeordinanceguidelines.pdf

<http://www.ufei.calpoly.edu/files/pubs/guidelines.pdf>

<https://www.arboday.org/programs/treeCityUSA/documents/sample-tree-ordinance.pdf>

<http://conservationtools.org/guides/37>

Urban Tree Ordinance Development Workbook: A preliminary guide book designed for communities developing new, or revising older, existing ordinances governing urban tree resources: The questionnaire / checklist: is attachment C.

http://conservationtools-production.s3.amazonaws.com/library_item_files/591/541/Urban_Tree_Ord_Development.pdf?AWSAccessKeyId=AKIAIQFJLILYGVDR4AMQ&Expires=1507222361&Signature=xGOVPbzo4FaVixjDwEJwgMFko1Q%3D

Print

Chico, CA Code of Ordinances

Chapter 16.68

VOLUNTARY HERITAGE TREE PROGRAM

Section:

16.68.010 Voluntary Heritage Tree Program.

16.68.010 Voluntary Heritage Tree Program.

A. The purpose and intent of the Voluntary Heritage Tree Program is to identify, promote public awareness of, maintain, and protect designated trees within the City of Chico. This program acknowledges that Heritage Trees, whether located on public or private property, are distinct and unique living resources of the City of Chico.

B. Any person may submit an application on a form supplied by the director and accompanied by the required fee, to designate a tree as a Heritage Tree. If an application is filed by a person other than the owner of the property on which the tree is located, then the application shall include the written and signed consent of the property owner and the property owner shall have the opportunity to be fully involved in the designation process. Applications shall be evaluated by the urban forest manager and forwarded to the Bidwell Park and Playground Commission for review and recommendation to the City Council. The City Council may designate a tree as a Heritage Tree if it meets any of the following criteria:

1. Any native Oak (*Quercus*) species or Sycamore (*Platanus*) species, having a diameter at breast height of thirty-six (36) inches or greater when a single trunk, or a cumulative diameter of thirty-six (36) inches or greater when a multi-trunk, and with good health and structure; or
2. The tree is an outstanding specimen of a desirable species of good health and quality structure; or
3. The tree is of historical interest; or
4. The tree is an unusual species, is of distinctive form, is a part of a significant grove or is otherwise unique.

C. Once a tree is designated as a Heritage Tree, a Notice of Heritage Tree Designation shall be recorded against the property on which the tree is located. Heritage Trees located on public property are exempt from the recording requirement.

D. Heritage trees may only be removed pursuant to Chapter 16.66 of this code.

(Ord. 2402)



GENERAL SERVICES DEPARTMENT

965 Fir Street
P.O. Box 3420
Chico, CA 95927-3420

(530) 896-7800
Fax (530) 895-4731
<http://www.ci.chico.ca.us>

CITY OF CHICO

APPLICATION FOR HERITAGE TREE

Property Owner: _____ Applicant (if different): _____

Owner's Name: _____ Applicant's Title: _____

Owner's Address: _____

Owner's Phone Number: _____ Applicant's Phone Number: _____

TREE INFORMATION

Location of tree(s): _____ (Address)

Diameter of tree(s): _____ Species of tree(s): _____

Tree Description and Reason for Request: (Enclose a map showing the exact location of the tree(s) and any proposed improvements along with any other documents which will help explain your request)

HAS THE TREE BEEN INSPECTED BY AN ARBORIST?

Tree Service or Arborist: _____ Phone #: _____

Assessment of Tree:

Date Signature of Property Owner

Date Signature of Applicant (If different from Owner)

\$150 Fee Enclosed? _____ Yes _____ No

CITY OF CHICO
HERITAGE TREE APPLICATION

TO:

DATE:

Your application requesting Heritage Tree Status

at (location of tree) _____

has been Inspected by the Urban Forest Manager and Found to be:

_____ An Oak or Sycamore 36" or larger

_____ An Outstanding Specimen of a Desirable Species in Good Health and Structure

_____ Of Historical Interest

_____ An Unusual Species, of distinctive form, part of a significant grove, or otherwise unique

_____ Other Information:

_____ Approved and will be forwarded to the Bidwell Park and Playground Commission

on: _____

_____ Denied for the following Reason(s):

(Date)

Urban Forester or authorized City Representative

BPPC Finding _____ Approved _____ Denied

Deed Recorded on _____ Date


Distribution: (When fully executed)

___ GSD ___ BO ___ ADPW-ENGR ___ PL DIR ___ CM

CHICO'S HERITAGE TREE PROGRAM

In March of 2010, after two years of sub-committee meetings and review, the Chico City Council approved several revisions in the Municipal Code relating to trees. This revision included setting up a Voluntary Heritage Tree Program (CMC Section 16.68.010) to recognize significant trees growing in Chico. On November 2, the City Council approved the first 6 Heritage trees. Three of those trees are located in Chico's Parks and Open Space. Another is a City street tree, and the final one was a private tree in a new development across from the Nature Center. Subsequently, 2 trees were nominated by CARD and approved for Heritage Tree Status on 12/21/10. the first private tree, a valley oak located on East 8th St, was nominated and approved in April, 2012. A street tree was nominated by its adjacent homeowner in January, 2013. These trees are described below. The City's program is active and seeking new Heritage Tree Applications.

A 67" valley oak (*Quercus lobata*) tree in Annie's Glen was nominated because of its prominent location next to the bike path near the entrance from Orient and E 2nd Street. The tree has a history of old flood damage at its base, where an old injury can be seen. Floods deposit soil over the original grade of the root collar, and sometimes, this soil can create a crown rot injury. The tree's root collar has been exposed, however, and the tree pruned to improve its safety and longevity.



This **53" diameter Blue Oak (*Quercus douglasii*)** is clearly seen by looking east from Baroni Neighborhood Park, and is located in the open space preserve in the Hillview Terrace Subdivision.. Blue oaks are found in the more upland areas of Chico, where the water table drops below 30' in depth. They therefore rely on an extensive root system to gather water for the long summer months, rather than the deep sinker roots of the valley oak. This makes them slower growing and more sensitive to construction that encroaches upon their canopy and root zone.

A mid-west species that was likely planted by the Bidwell's in Children's Playground, this **52" Sweetgum (*Liquidambar styraciflua*)** is one of the largest in California – rivaling one in Sacramento that is considered the largest at 54". Our tree is an excellent specimen, with little to no limb breakage and is found at the north end of the Park. It is usually in full fall color for most of November and December.





The city's largest Heritage Tree is a 78" Valley oak located near the corner of Eaton and Cohasset and a 66" oak on Heritage Oak Drive. This tree grows on the bank of Sycamore Creek. At some point in time in the distant past, the soil was raised over the root collar of the tree, probably in the building of the levy and Eaton Road, to the south of the tree. Volunteers have cared for the tree over the years by installing large logs around the outside of the drip line to keep vehicles out from under the canopy, pruning and cabling the tree, and removing the excess soil over the root collar. Called the Eaton Oak, it is an incredible specimen that is a joy to stand under. One limb has grown to the ground and is helping to support the massive structure.

A tree that the City can be proud of preserving is found on Heritage Oak Drive. This 67" valley oak was recognized by the developer and care was taken to keep house foundations at the outer edge of the canopy. The street was carefully constructed around the tree as well. Not long after the tree was preserved, it broke a large limb growing to the north, so now the tree's canopy is pruned on a regular basis to reduce the weight on the long horizontal limbs so they don't break. An area at the base of the tree is mulched on a regular basis, and was made part of the street's right of way for the tree.




A 70" Bastogne or Paradox walnut (*Juglans hindsii x regia*) located at the new Park Forest Subdivision is a seedling hybrid between the English and the native black walnut. Fairly common in the Chico area, this tree was planted in 1900 as a pollinator for a small family owned walnut orchard. Property developers of Park Forest LLC set aside the area around the tree to protect it and feature the tree as a center piece of their Park Forest project on East 8th Street across from the Nature Center. They built a large ring around the dripline to protect it during construction. To date no significant damage has occurred from construction.

Two additional trees – located in the CARD Community Park off 20th Street were nominated for the program. On the south side of the park, at the end of Cleveland Street, a lovely 62" valley oak provides shade and beauty for a small picnic area. The tree existed long before the park was built, and has been pruned and cabled over the years.



A California black walnut (*Juglans hindsii*), of the type that John Bidwell planted around Chico, covers another picnic area north of the oak. It sits near the children's playground in the 20th Street Community Park. These trees are known for their exceedingly beautiful wood that is also quite valuable. The nuts are excellent eating, although cracking the nut is so difficult that they are no longer produced in California. They remain an agricultural necessity, though, and are used for root stock on most walnuts because they are resistant to the oak root fungus disease common to California soils.

A privately owned valley oak located at 2135 East 8th Street was nominated by the property owners in February of 2012. The tree is in excellent condition and provides both shade to the home. It also is a main focal point of the yard, and can easily be seen from East 8th Street.





Blue atlas cedars are planted throughout California, but rarely does one attain the size and grace of the tree located on the side of 1641 Mulberry. This tree has grown to over 50" in diameter, despite having been struck by lightning at some point in the past. The injury can still be seen on the central trunk, which is one reason why the two side branches have become dominant. This cedar is a City street tree.

HELP US FIND MORE HERITAGE TREES!

Citizens owning significant trees who are interested in having them considered for Heritage Tree Status are encouraged to contact the City's Tree Division at (530) 896-7800, or email parkinfo@Chicoca.gov



BPPC Tree Committee Report

Meeting Date 10/12/2017

DATE: 10/12/17
TO: Bidwell Park and Playground Commission (BPPC) Tree Committee
FROM: Richie Bamlet, Urban Forest Manager
SUBJECT: Consideration of educational topics to the public in an effort to protect, preserve and enhance the City of Chico's street trees and urban forest.

RECOMMENDATION

Staff seeks continued discussion from the Committee on proposed educational topics in an effort to protect, preserve and enhance the City of Chico's street trees and urban forest.

BACKGROUND:

At the previous Tree Committee meeting on 8/10/17 possible outreach and education efforts were discussed. Committee members returned with six draft topic headers which are presented for discussion. Staff seeks to develop these topics into actionable items that can be delivered as educational products to the citizens of Chico.

DISCUSSION:

Committee member Haar provided the following draft topics for a: 12-month Tree Public Education Program to be presented on a bi-monthly basis:

Overview:

To provide educational topics to the public in an effort to protect preserve and enhance the City of Chico Street Trees and City's Urban Forest

1. City Permits and Restrictions/Fines : Removal Request and Appeals
2. Right Tree Right Place: where and how to plant
3. Proper pruning for structure and form: safety 7' sidewalk 14' road. Root pruning safety
4. Heritage Oak History and Permits.
5. Benefits of a healthy Urban Forest and Tree Canopy: City Tree watering
6. Sudden Limb drop

FISCAL IMPACT:

None. Grant assistance might be available for implementation.